

MULTIPOLE ANALYSIS OF RECENT PION ELECTROPRODUCTION DATA

R. ARNDT, W. BRISCOE, I. STRAKOVSKY, AND R. WORKMAN

*Center for Nuclear Studies, Department of Physics
The George Washington University, Washington, D.C. 20052*

We have performed a series of global and single- Q^2 fits to the growing database for pion electroproduction. We compare to some other recent fits which have used different formalisms, more selective datasets, and truncated multipole expansions. We present results for the E2/M1 and S2/M1 ratios, as a function of Q^2 .

1. Fit Procedure

In fitting the pion electroproduction database ¹, we have fixed the $Q^2 = 0$ point based on our fits to pion photoproduction ². The photoproduction multipoles have been parametrized using the form

$$M = (\text{Born} + \alpha_B)(1 + iT_{\pi N}) + \alpha_R T_{\pi N} + (\text{Im}T_{\pi N} - T_{\pi N}^2)(\alpha_r^H + i\alpha_i^H), \quad (1)$$

containing the Born terms, with phenomenological pieces (α) maintaining the correct threshold behavior and Watson's theorem. The πN T-matrix ($T_{\pi N}$) connects each multipole to structure found in the elastic πN scattering analysis. Additional structure is parametrized through the Born terms (which contain vector-meson contributions), and the phenomenological structure functions (α_B , α_R , α_r^H , and α_i^H).

At non-zero Q^2 , the Born terms have built-in Q^2 dependence, other terms have been modified by a factor

$$f(Q^2) = \frac{k}{k(Q^2=0)} \frac{1}{(1 + Q^2/0.71)^2} e^{-\Lambda Q^2} \left(1 + \alpha Q^2 + \beta \left[\frac{W}{W_R} - 1 \right] \right), \quad (2)$$

where k is the photon CM momentum, Λ is a universal cutoff factor, and the constants, α and β , are searched for each multipole. We have performed energy and Q^2 -dependent fits over the full kinematic range [W to 1.7 GeV and Q^2 to 5 (GeV/c)²] in addition to fitting data clustered around particular Q^2 values.




Figure 1. Energy-angle distribution of $\pi^0 p$ and $\pi^+ n$ electroproduction data: $Q^2 = 0.40, 0.50, 0.65, 0.75, 0.90, 1.15, 1.45$, and $1.80 (\text{GeV}/c)^2$ associated with JLab CLAS $\pi^0 p$ Q^2 binning for single- Q^2 results.




Figure 2. $E2/M1$ and $S2/M1$ ratios *vs* Q^2 . Values were extracted from our fixed Q^2 analyses starting on the global fit (filled squares) using world and JLab CLAS data³. Results from Ref.³ (filled circles) are given. In addition, (left panel), our pion photoproduction result ($Q^2 = 0$)² (filled asterisk), and (right panel), the data of Refs.⁴ (open triangle) and⁵ (open diamond) are shown. The solid curves give our global (energy and Q^2 dependent) best-fit results. Dash-dotted and dashed curves are from Refs.⁶ and⁷, respectively.




Figure 3. Total cross section for $\pi^0 p$ electroproduction, showing the variation with Q^2 .

As can be seen in Figure 1, there is relatively little $\pi^+ n$ data at moderate Q^2 . As a result, some analyses actually quote a ratio of $E2/M1$ for the $\pi^0 p$ charge channel, rather than the isospin 3/2 component, assuming a negligible isospin 1/2 part at the Δ -resonance.

1.1. Data Reduction

A fit to distributions in θ and ϕ can be made more efficient if the symmetry $\sigma(\phi) = \sigma(360 - \phi)$ is used. Let $\sigma_1 = \sigma(\phi)$, $\sigma_2 = \sigma(360 - \phi)$, and z be the statistical error (squared). If we take σ to be the fitted value, then for each




Figure 4. Differential cross section for $\pi^0 p$ electroproduction, showing the ϕ dependence. Data are from ³.




Figure 5. Q^2 dependence of the magnetic (M_{1+}), electric (E_{1+}), and scalar (S_{1+}) multipoles, for isospin 3/2.

pair of ϕ values we have

$$\chi^2 = \frac{(\sigma - \sigma_1)^2}{z_1} + \frac{(\sigma - \sigma_2)^2}{z_2}, \quad (3)$$

which can be written as

$$\chi^2 = \frac{(\sigma_1 - \sigma_2)^2}{z_1 + z_2} + \frac{(\sigma - \sigma_a)^2}{z_a}, \quad (4)$$

where $\sigma_a = (z_1\sigma_2 + z_2\sigma_1)/(z_1 + z_2)$ is the weighted average of σ_1 and σ_2 , and z_a is given by

$$\frac{1}{z_a} = \frac{1}{z_1} + \frac{1}{z_2}. \quad (5)$$

Using the form given in Eq. (4), only half of the data points are required in the fitting routine, the first term being a constant which measures the internal consistency of a given set. For the set of $\pi^0 p$ data given in Ref. ³, for example, the first term in Eq. (4) contributes a χ^2 of 15040 for the 12528 ϕ pairs.

2. Results

In Figures 2–5, we compare our result to MAID ⁶, Sato-Lee ⁷, and single- Q^2 fits performed by the Jefferson Lab CLAS ³ Collaboration. Below 2 (GeV/c)², there is considerable scatter, with SAID and MAID extrapolating to an E2/M1 zero-crossing; no crossing being seen in Sato-Lee. As indicated above, the higher- Q^2 region is sparsely populated. In our fits, we find that the inclusion of additional preliminary data can shift the crossing point significantly, but a crossing always occurs. This is in particular disagreement with the 4 (GeV/c)² Hall C value ⁸, and details of the various fits, at this kinematic point, are being studied in more detail.

Acknowledgments

This work was supported in part by a U. S. Department of Energy Grant No. DE-FG02-99ER41110. We also acknowledge a contract from Jefferson Lab under which part of this work was done. Jefferson Lab is operated by the Southeastern Universities Research Association under DOE contract DE-AC05-84ER40150.

References

1. R. A. Arndt, I. I. Strakovsky, and R. L. Workman, *πN Newsletter*, 16, 150 (2002).
2. R. A. Arndt, W. J. Briscoe, I. I. Strakovsky, and R. L. Workman, *Phys. Rev. C* **66**, 055213 (2002).
3. K. Joo *et al.* (The CLAS Collaboration), *Phys. Rev. Lett.* **88**, 122001 (2002). We restrict our fits to bins having more than 5 events per bin.
4. Th. Pospischil *et al.*, *Phys. Rev. Lett.* **86**, 2959 (2001).
5. F. Kalleicher *et al.*, *Z. Phys.* **A359**, 201 (1997).

6. The extended version of the MAID analysis refers to DMT2001 <http://www.kph.uni-mainz.de/MAID/>. See also S. S. Kamalov, S. N. Yang, D. Drechsel, O. Hanstein, and L. Tiator, *Phys. Rev. C* **64**, 032201 (2001); D. Drechsel, O. Hanstein, S. S. Kamalov, and L. Tiator, *Nucl. Phys.* **A645**, 145 (1999).
7. T. Sato and T. S. H. Lee, *Phys. Rev. C* **63**, 055201 (2001).
8. V. V. Frolov *et al.* *Phys. Rev. Lett.* **82**, 45 (1999).